



Colorado State University Veterinary Medical Center Feline Acute Pain Scale



Pain Score	Example	Psychological & Behavioral	Response to Palpation	Body Tension
No Score		<input type="checkbox"/> Animal is sleeping and cannot be evaluated		
0		<input type="checkbox"/> Content and quiet when unattended <input type="checkbox"/> Comfortable when resting <input type="checkbox"/> Interested in or curious about surroundings	<input type="checkbox"/> Not bothered by palpation of wound or surgery site, or to palpation elsewhere	Minimal
1		<input type="checkbox"/> Signs are often subtle and not easily detected in the hospital setting; more likely to be detected by the owner(s) at home <input type="checkbox"/> Earliest signs at home may be <u>withdrawal from surroundings or change in normal routine</u> <input type="checkbox"/> In the hospital, may be content or slightly unsettled <input type="checkbox"/> Less interested in surroundings but will look around to see what is going on	<input type="checkbox"/> May or may not react to palpation of wound or surgery site	Mild
2		<input type="checkbox"/> Decreased responsiveness, seeks solitude <input type="checkbox"/> Quiet, loss of brightness in eyes <input type="checkbox"/> Lays curled up or sits tucked up (all four feet under body, shoulders hunched, head held slightly lower than shoulders, tail curled tightly around body) with eyes partially or mostly closed <input type="checkbox"/> Hair coat appears rough or fluffed up <input type="checkbox"/> May intensively groom an area that is painful or irritating <input type="checkbox"/> Decreased appetite, not interested in food	<input type="checkbox"/> Responds aggressively or tries to escape if painful area is palpated or approached <input type="checkbox"/> Tolerates attention, may even perk up when petted as long as painful area is avoided	Mild to Moderate Reassess analgesic plan
3		<input type="checkbox"/> Constantly yowling, growling, or hissing when unattended <input type="checkbox"/> May bite or chew at wound, but unlikely to move if left alone	<input type="checkbox"/> Growls or hisses at non-painful palpation (may be experiencing allodynia, wind-up, or fearful that pain could be made worse) <input type="checkbox"/> Reacts aggressively to palpation, adamantly pulls away to avoid any contact	Moderate Reassess analgesic plan
4		<input type="checkbox"/> Prostrate <input type="checkbox"/> Potentially unresponsive to or unaware of surroundings, difficult to distract from pain <input type="checkbox"/> Receptive to care (even mean or wild cats will be more tolerant of contact)	<input type="checkbox"/> May not respond to palpation <input type="checkbox"/> May be rigid to avoid painful movement	Moderate to Severe May be rigid to avoid painful movement Reassess analgesic plan



Comments _____